



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.



Letter to the Editor

Can COVID-19 cause sudden sensorineural hearing loss?



We thank Pietro et al. for their comments on our recently published article (Kilic et al., 2020). In previous studies, viral infections have been reported to have an important role in the etiology of sudden sensorineural hearing loss (SSNHL) (Cohen et al., 2014). Therefore, it is not unreasonable to think that coronavirus disease 2019 (COVID-19), as a viral infection, may also cause SSNHL. Moreover, studies on the effects of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) virus on central and peripheral nervous systems have been recently published. Some of these studies have reported Guillain-Barré syndrome (Toscano et al., 2020) and impaired olfactory function (Eliezer et al., 2020) to be related to SARS-CoV-2. Another recently published study reported neurologic changes in 58 of 64 patients with COVID-19 (Helms et al., 2020).

When we observed an unexpected increase in SSNHL cases with no other symptoms in our ENT outpatient clinics during the COVID-19 pandemic, we thought that this might be related to COVID-19. Therefore, we investigated the possibility of SARS-CoV-2 infection in these patients. Using RT-PCR, we detected the presence of SARS-CoV-2 in one of our five patients (Kilic et al., 2020). Our study is the first in the literature to show that there may be a relationship between SSNHL and COVID-19.

The results of a recent study on the effect of COVID-19 on hearing functions supports our hypothesis (Mustafa, 2020). The author reported that both transient evoked otoacoustic emissions and high frequency pure-tone threshold amplitudes were significantly worse in the asymptomatic COVID-19 PCR-positive patient group.

Undoubtedly, routine healthcare services are disrupted during this pandemic, so the association of SSNHL cases with COVID-19 may have gone unnoticed in healthcare institutions due to the increased workload. Of course, in order to make a definitive diagnosis, studies demonstrating the presence of the virus in the relevant tissues, such as postmortem studies, are required. However, in the ongoing pandemic, we believe that every symptom should be considered and shared with the medical community. We believe that our article will increase awareness of this issue. This will encourage the design of large studies, in regions with high prevalence of COVID-19, as also suggested by Pietro et al.

Conflict of interest

The authors declare that they have no conflicts of interest.

References

- Cohen BE, Durstenfeld A, Roehm PC. Viral causes of hearing loss: a review for hearing health professionals. *Trends Hear* 2014;18;. doi:<http://dx.doi.org/10.1177/2331216514541361>
- Eliezer M, Hautefort C, Hamel AL, Verillaud B, Herman P, Houdart E, et al. Sudden and complete olfactory loss function as a possible symptom of COVID-19. *JAMA Otolaryngol Head Neck Surg* 2020;146(7):674–5. doi:<http://dx.doi.org/10.1001/jamaoto.2020.0832>.
- Helms J, Kremer S, Merdji H, Clere-Jehl R, Schenck M, Kummerlen C, et al. Neurologic features in severe SARS-CoV-2 infection. *N Engl J Med* 2020;382(23):2268–70. doi:<http://dx.doi.org/10.1056/NEJMc2008597>.
- Kilic O, Kalcioğlu MT, Cag Y, Tuysuz O, Pektaş E, Caskurlu H, et al. Could sudden sensorineural hearing loss be the sole manifestation of COVID-19? An investigation into SARS-CoV-2 in the etiology of sudden sensorineural hearing loss. *Int J Infect Dis* 2020;97:208–11. doi:<http://dx.doi.org/10.1016/j.ijid.2020.06.023>.
- Mustafa MWM. Audiological profile of asymptomatic Covid-19 PCR-positive cases. *Am J Otolaryngol* 2020;41(3):102483. doi:<http://dx.doi.org/10.1016/j.amjoto.2020.102483>.
- Toscano G, Palmerini F, Ravaglia S, Ruiz L, Invernizzi P, Cuzzoni MG, et al. Guillain-Barré syndrome associated with SARS-CoV-2. *N Engl J Med* 2020;382(26):2574–6. doi:<http://dx.doi.org/10.1056/NEJMc2009191>.

Mahmut Tayyar Kalcioğlu*

Istanbul Medeniyet University, Faculty of Medicine, Department of Otorhinolaryngology, Istanbul, Turkey

Yasemin Cag

Istanbul Medeniyet University, Faculty of Medicine, Department of Infectious Diseases and Clinical Microbiology, Istanbul, Turkey

Osman Kilic

Ozan Tuysuz

Istanbul Medeniyet University, Faculty of Medicine, Department of Otorhinolaryngology, Istanbul, Turkey

* Corresponding author.

E-mail address: mahmut.kalcioğlu@madaniyet.edu.tr
(M. Kalcioğlu).

Received 24 September 2020